


**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

**APPROVALS**

	<u>Name</u>	<u>Date</u>	<u>Signature</u>
Original Document Author:	Dr Caroline Leech		
Revised Document Prepared by:	Dr Stuart Reid		
Reviewed by:	Danny Evans, Paramedic Phil Bridle, Head of Operations		
CGG Approval:	Dr Justin Squires, Deputy Clinical Lead		
Next Review Date:	March 2021		

**HISTORY**

<b>Effective Date</b>	<b>Version No.</b>	<b>Summary of Amendment</b>
14/12/12	2.0	Reviewed, no changes, reformatted into new format
Feb 2014	2.1	Updates
Oct 2017	2.2	Review
24/11/17	2.3	Update of advice re patients with neck pain on page 4.
Feb 2020	2.4	Reviewed, no amendments required.

**APPENDICES**

<b>Document Reference Number</b>	<b>Document Title</b>
Appendix A	Canadian C Spine rules algorithm
Appendix B	JRCALC Spinal Immobilisation algorithm



## Title CSOP 009                      Spinal Injury

Version No: 2.4

Effective date: 16/03/2020

### 1. Purpose / Scope

This CSOP provides guidance on spinal immobilisation and the management of patients with suspected spinal cord injury.

### 2. Definitions/Acronyms:

Abbreviations	Definitions
CSOP	Clinical Standard Operating Procedure
JRCALC	Joint Royal College Ambulance Liaison Committee
RSI	Rapid Sequence Induction
TAAS	The Air Ambulance Service
PRF	Patient Report Form
TARN	Trauma Audit Research Network
NICE	National Institute for Health and Care Excellence

### 3. Introduction

Spinal cord injury after trauma is described in around 2% of patients on the TARN database, and data from Victoria, Australia gives a figure of 0.5% of patients who are immobilised pre hospital. Driven by traditional teaching and dogma around the need for cervical spine immobilisation, and perceived medicolegal concerns, UK practice around immobilisation has become extraordinarily conservative.

### 4. 'Clearing' the Spine

JRCALC clinical practice guidelines 2016 describe an algorithm for the prehospital removal of spinal immobilisation and this has been reproduced in Annex A. This algorithm is based upon the NEXUS low risk criteria, a clinical decision rule used to assess the need for cervical spine radiography in Emergency Departments.

NICE Spinal Injury guidelines 2016 (1) advocate the use of the Canadian C Spine rules in the pre hospital setting, and the Canadian C Spine rules have been validated for use by paramedics (2). **In line with NICE, it is acceptable for TAAS clinical staff to apply the Canadian C Spine rule in their TAAS practice (appendix for algorithm), allowing C Spines to be 'cleared' at scene.**



**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

NICE acknowledges that the use of the Canadian C spine rules in children may be difficult and that their developmental age should be considered. Furthermore current APLS guidance has de-emphasised the use of cervical collars in children.

The concept of a 'distracting injury' invalidating the clinical assessment of the cervical spine has been disproved in the literature and should not routinely influence practice. (3)

**Spinal Immobilisation**

The perceived aim of spinal immobilisation is to reduce the risk of causing, or worsening, spinal cord injury by the movement of unstable spinal fractures. Precautions taken to promote immobilisation include:

- Manual in-line stabilisation
- The application of a rigid cervical collar
- A transfer / transport device that maintains the patient in-line (e.g. scoop stretcher or vacuum mattress)
- Head blocks and tape
- Patient handling with the aim of maintaining the spine in-line

However, spinal immobilisation is not a benign intervention; it can cause:

- Pain or discomfort
- Obstruction to cerebral venous return
- Airway compromise
- Increased aspiration risk
- Pressure areas
- Reduced ventilatory ability
- Catastrophic deterioration in patients with a fixed spinal deformity such as ankylosing spondylitis
- Distraction of personnel on scene from life threatening issues

Furthermore, there is no high quality evidence to suggest that this practice reduces secondary cord injury amongst the small proportion of trauma patients with spinal injury. (4,5,6)

Clinicians should therefore consider the possibility of spinal injury in every trauma patient and make a dynamic assessment as to the appropriateness of spinal precautions for individual patients.



**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

---

The following offers guidance as to appropriate practice by TAAS crews:

- 1. Alert and cooperative patients with no neck symptoms – no indication for immobilisation**
- 2. Alert and cooperative patients with neck symptoms or alert and cooperative patients with clear neurological signs of spinal cord injury - gentle manual in line stabilisation during primary survey. Blocks for transfer on scoop stretcher.**
- 3. Agitated / confused / decreased GCS patients – no indication for immobilisation as liable to increase agitation and confusion, put cervical spine under more strain and expose patient to more risks from side effects of immobilisation**
- 4. Anaesthetised or sedated patients where neck injury is suspected – gentle manual in line stabilisation during primary survey and airway interventions (as per NICE). Blocks for transfer on scoop stretcher.**

Unnecessary movements of the spine should be avoided in all patients in the pre-hospital phase.

If a decision is made to apply spinal precautions, this should not unduly delay the primary survey or impede the delivery of time-critical interventions. This includes the rapid extrication of a time critical patient, for example from a vehicle.

If treating a child with a suspected spinal injury, a pragmatic approach is required. This may include keeping an infant in a car seat, use of rolled blankets, vacuum mattresses or splints, and family members to assist. (NICE 2016)

There is evidence of harm by immobilising the cervical spine in patients with penetrating neck injuries. This is unnecessary and should be avoided.



**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

**5. Extrication**

When there is immediate threat to a person's life and rapid extrication is needed, make all efforts to limit spinal movement without delaying treatment.

In line with the consensus statement from the Faculty of Prehospital Care (7), it is recommended that alert and cooperative patients who are not trapped should be invited to extricate themselves from a vehicle and lie on a stretcher to allow assessment. If proceeding with self-extrication, it should be explained to the patient that if they develop any spinal pain, numbness, tingling or weakness, they should stop moving and wait to be moved.

Do not transport people with suspected spinal injury on a longboard or any other extrication device. A longboard should only be used as an extrication device.

**6. Documentation and handover**

When TAAS clinicians have determined that spinal precautions are not necessary this should be clearly documented on the patient report form, in particular the distinction between a non-immobilised, but not cleared neck; versus a non-immobilised, cleared neck.

Be aware that ambulance crews working to JRCALC may find it difficult to hand over patients to a hospital where a decision has been made by a TAAS clinician at the scene. It should also be noted that the JRCALC clinical practice guidelines followed by personnel from tasking ambulance services mandate that when spinal precautions are applied these should be 'full', i.e. the whole spine should be immobilised not just the neck or back. If TAAS clinicians have determined that selectively immobilising only part of the spine is clinically appropriate then they should clearly document this in the patient report form.



**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

**Spinal Cord Injury – Clinical Presentation**

Spinal cord injury may present with the following symptoms and signs in conscious patients:

- Neck or back pain
- Loss of movement or sensation in limbs
- Sensation of burning or electric shocks may be experienced in the trunk or limbs.
- Breathing does not feel normal
- Patient may not experience pain even from significant injury
- Diaphragmatic or abdominal breathing
- Hypotension with bradycardia
- Warm peripheries and vasodilatation despite having a low blood pressure
- Flaccid muscles with absent reflexes
- Partial or complete priapism

**7. Assessment of Spinal Cord Injury**

As part of the disability assessment for all trauma patients, movement of all four limbs should be assessed and documented (especially prior to any sedation or RSI). Remember that your documentation may be essential for clinical or legal purposes at a later date.

Where a spinal fracture or cord injury is suspected the assessment should consider if the patient:

- has any spinal pain
- has any hand or foot weakness (motor assessment)
- has altered or absent sensation in the hands or feet (sensory assessment)
- has priapism (unconscious or exposed male)
- has a history of past spinal problems, including previous spinal surgery or conditions that predispose to instability of the spine.

(NICE 2016)


**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

**Suspected thoracic and lumbar injuries**

Where any of the factors below are present, NICE recommend full spinal immobilisation with a scoop stretcher. This is the recommended approach at TAAS, mostly as it will facilitate an easy transfer to the Emergency Department trolley with minimal patient movement. It is acceptable for TAAS clinicians not to extend this immobilisation to the neck, instead managing the patient according to the above guidance.

- age 65 years or older and reported pain in the thoracic or lumbosacral spine
- dangerous mechanism of injury (fall from a height of greater than 3 metres, axial load to the head or base of the spine – for example falls landing on feet or buttocks, high-speed motor vehicle collision, rollover motor accident, lap belt restraint only, ejection from a motor vehicle, accident involving motorised recreational vehicles, bicycle collision, horse riding accidents)
- pre-existing spinal pathology, or known or at risk of osteoporosis – e.g. steroid use
- suspected spinal fracture in another region of the spine
- abnormal neurological symptoms (paraesthesia or weakness or numbness)
- on examination:
  - abnormal neurological signs (motor or sensory deficit)
  - new deformity or bony midline tenderness (on palpation)
  - bony midline tenderness (on percussion)
  - midline or spinal pain (on coughing)
- pain or abnormal neurological symptoms on mobilisation

This should all be clearly recorded on the PRF and communicated in the handover at hospital.



**Title CSOP 009                      Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

**8. Treatment of Spinal Cord Injury (SCI)**

Several clinical series of human patients with acute SCI managed in an aggressive fashion with attention to blood pressure, oxygenation, and hemodynamic performance report no deleterious effects of therapy and suggest improved neurological outcome. (8)

**A**

Consider intubation for ventilation if the patient has respiratory failure, bearing in mind that SCI patients can be extremely difficult to wean later in their clinical course.

**B**

The spinal cord can suffer secondary injury (in the same manner as a head injury) and therefore hypoxia must be avoided.

Ask the patient if their breathing feels normal and observe for diaphragmatic breathing as this may indicate a high cervical spine injury. Check the SaO<sub>2</sub>, and check the EtCO<sub>2</sub> of the non-intubated patient- an abnormally high or increasing level may indicate impending ventilatory failure.

**C**

Hypotension must be corrected. High quality evidence is lacking in terms of a target blood pressure, although many authors advocate a mean arterial blood pressure of 85-90mmHg (8,9). Vasopressors should be used early and consideration given to treating a symptomatic bradycardia. (useful drugs include ephedrine, adrenaline or metaraminol). Excess fluid administration can lead to morbidity.

In the polytrauma patient, look for causes of hypotension, e.g. tension pneumothorax, hypovolaemia, tamponade, etc. and treat as per standard procedures. When all other causes have been excluded or treated and the patient is thought to be euvoalaemic it may be appropriate to initiate inotropic support.

**D**

Ensure a full neurological examination has been performed and documented. Pay close attention to patient handling with the aim of preventing further neurological injury. Spinal-injured patients are at high risk of pressure sores from an early stage.

**E**

Temperature control - patients with spinal cord injury loses the ability to thermo-regulate and therefore must be kept warm.



**Title CSOP 009 Spinal Injury**

Version No: 2.4

Effective date: 16/03/2020

## 9. Triage and transfer – spinal cord injury

As with any other Major Trauma patient the regional Major Trauma Triage Tool should be applied. Do not transfer patients with suspected spinal cord injury direct to the regional Spinal Injuries unit. These patients should be triaged to a Major Trauma Centre. Remember that paralysis may mask other traumatic injuries.

### **FURTHER READING / REFERENCES**

- 1) **NICE Guidance [NG41] Spinal injury: assessment and initial management**  
**Published date: February 2016**
- 2) **Vaillancourt et al. The out-of-hospital validation of the Canadian C-Spine Rule by paramedics. Ann Emerg Med. 2010 Jan;55(1):22.**
- 3) **Rose et al. Clinical clearance of the cervical spine in patients with distracting injuries: It is time to dispel the myth. J Trauma Acute Care Surg 2012.Aug;73(2):498-502.**
- 4) **Sundstrom et al. Prehospital Use of Cervical Collars in Trauma Patients: A Critical Review. J Neurotrauma. 2014 Mar 15;31(6):531-40**
- 5) **Oteir et al. Should suspected cervical spinal cord injury be immobilised?: A systematic review. Injury 2015.46:528–535**
- 6) **Oto et al. Early Secondary Neurologic Deterioration After Blunt Spinal Trauma: A Review of the Literature. Acad Emerg Med 2015;22:1200–1212**
- 7) **Connor et al. Pre-hospital spinal immobilisation: an initial consensus statement. Emerg Med J 2013;30:1067-1069**
- 8) **Ryken et al. The Acute Cardiopulmonary Management of Patients With Cervical Spinal Cord Injuries. Neurosurgery 72:84–92, 2013**
- 9) **Walters et al. Guidelines for the Management of Acute Cervical Spine and Spinal Cord Injuries: 2013 Update. Neurosurgery 2013;60 S1:82-91**

**End of Document**